



UNIVERSIDADE
ESTADUAL DE LONDRINA

FABIANE FERREIRA MONTEIRO

**OBESIDADE E ATIVIDADE FÍSICA NA VIDA DIÁRIA EM
PACIENTES COM DPOC**

FABIANE FERREIRA MONTEIRO

**OBESIDADE E ATIVIDADE FÍSICA NA VIDA DIÁRIA EM
PACIENTES COM DPOC**

Dissertação apresentada ao Programa de Pós-Graduação em Ciências da Reabilitação (Programa Associado entre Universidade Estadual de Londrina [UEL] e Universidade Norte do Paraná [UNOPAR]), como requisito parcial à obtenção do título de Mestre em Ciências da Reabilitação.

Orientador: Prof. Dr. Fábio de Oliveira Pitta

Londrina
2011

**Catálogo elaborado pela Divisão de Processos Técnicos da Biblioteca Central da
Universidade Estadual de Londrina**

Dados Internacionais de Catalogação-na-Publicação (CIP)

M775o Monteiro, Fabiane Ferreira.
Obesidade e atividade física na vida diária em pacientes com DPOC /
Fabiane Ferreira Monteiro. – Londrina, 2011.
43 f.

Orientador: Fábio de Oliveira Pitta.
Dissertação (Mestrado em Ciências da Reabilitação) – Universidade
Estadual de Londrina, Centro de de Ciências da Saúde, Programa de
Pós-Graduação em Ciências da Reabilitação, 2011.
Inclui bibliografia

1. Pulmões – Doenças – Exercícios físicos – Teses. 2. Pulmões –
Doenças – Reabilitação – Teses. 3. Obesidade – Exercícios físicos – Teses.
I. Pitta, Fábio de Oliveira. II. Universidade Estadual de Londrina. Centro de
Ciências da Saúde. Programa de Pós-Graduação em Ciências da
Reabilitação. III. Título.

CDU 615.8:616.24

FABIANE FERREIRA MONTEIRO

**OBESIDADE E ATIVIDADE FÍSICA NA VIDA DIÁRIA EM PACIENTES
COM DPOC**

Dissertação apresentada ao Programa de Pós-Graduação em Ciências da Reabilitação (Programa Associado entre Universidade Estadual de Londrina [UEL] e Universidade Norte do Paraná [UNOPAR]), como requisito parcial à obtenção do título de Mestre em Ciências da Reabilitação.

BANCA EXAMINADORA

Orientador. Prof. Dr. Fábio de O. Pitta
UEL – Londrina – PR

Componente da banca. Prof.^a Dr.^a Vanessa S.
Probst
UNOPAR – Londrina – PR

Componente da banca. Prof.^a Dr.^a. Irma de
Godoy
UNESP – São José do Rio Preto – SP

Londrina, 09 de dezembro de 2011

AGRADECIMENTOS

Gostaria de iniciar agradecendo a Deus, que neste assim como em todos os outros desafios por mim enfrentados esteve sempre ao meu lado me dando força, sabedoria e coragem.

Minha imensa gratidão é dedicada ao meu orientador Prof. Dr. Fábio Pitta. Graças a você meu sonho pode se tornar realidade. Você me ensinou, me ajudou e me apoiou em todos os momentos em que precisei. As suas conquistas são simplesmente o reflexo da sua paixão e dedicação pelo que faz. Apesar de todos os seus compromissos, que são MUITOS, você sempre consegue fazer com que todos que estão a sua volta sintam-se pessoas importantes. Você é uma inspiração para todos nós que temos o privilégio de conviver com você. Te admiro como profissional e pesquisador e principalmente como o ser humano de Deus que você é. Muito obrigada por tudo! Espero ter a oportunidade de aprender muito mais com você.

Sou muito grata aos meu pais, Edson e Lucimarci, que mesmo nos momentos mais difíceis sempre colocaram minha educação em primeiro lugar. Agradeço por todo apoio, por todo carinho e todo amor que sempre dedicaram a mim. Sei o quanto se orgulham de mim e dedico tudo que conquisto a vocês! Agradeço também ao meu irmão Felipe, que na correria do dia-a-dia pouco tempo tenho tido para conversarmos sobre suas novidades (que são muitas), e a minha querida avó Benedita que faz tudo por mim. Amo vocês!

Agradeço ainda ao meu noivo, Gustavo, que sempre compreendeu minhas viagens para congressos, minha falta de tempo e até mesmo meu mau humor pelo excesso de trabalho. Você é meu presente de Deus! Seu amor e carinho me tornam uma pessoa melhor.

A todos os professores deste programa de mestrado, que contribuíram para meu crescimento profissional e científico, meu muito obrigada!

Agradeço também aos professores da banca, Dra. Irma Godoy, que prontamente atendeu ao nosso convite e com certeza contribuirá muito com minha pesquisa, e Dra Vanessa Probst que não só neste momento mas em muitos outros já contribuiu muito para meu crescimento.

Meu sincero muito obrigada ainda a todos os colegas do LFIP, alunos e agregados, que foram fundamentais para que esse trabalho se realizasse.

Por fim, agradeço todos aqueles que eventualmente aqui não foram citados mas que de alguma forma foram importantes para que este sonho fosse realizado.

MONTEIRO, Fabiane. **Obesidade e atividade física na vida diária em pacientes com DPOC**. 2011. 43 f. Trabalho de Conclusão de Curso do Programa de Pós-Graduação em Ciências da Reabilitação (Programa Associado entre Universidade Estadual de Londrina [UEL] e Universidade Norte do Paraná [UNOPAR]) – Universidade Estadual de Londrina, Londrina, 2011.

RESUMO

Objetivo: Estudar as diferenças na composição corporal de pacientes com doença pulmonar obstrutiva crônica (DPOC) fisicamente ativos e inativos; e verificar se existe relação entre obesidade e atividade física na vida diária nesta população. **Métodos:** Composição corporal (bioimpedância elétrica) e nível de atividade física na vida diária (AFVD) (monitores de atividade física: DynaPort e SenseWear) foram avaliados em 74 pacientes com DPOC (45 homens, 65±9anos, VEF₁=40±15%pred, IMC=27±6kg.m⁻²). Os pacientes foram inicialmente separados em 2 grupos: fisicamente ativos (> 30 minutos/dia de tempo em atividade física [TAF] de intensidade no mínimo moderada, ou TAF>moderada) e inativos (não atingiam essas recomendações). A mesma amostra foi ainda dividida com base no índice de massa corpórea (IMC) em subnutridos, eutróficos, sobrepeso e obesos; e com base no percentual de gordura corporal (%GC) em obesidade moderada, obesidade elevada e obesidade mórbida. **Resultados:** No grupo completo, TAF>moderada se correlacionou modestamente com IMC (r=-0.28; p=0.02) e com a GC (Kg) (r=-0.30; p=0.001), mas não com a massa magra corporal (MMC) (Kg) (r=-0.18; p=0.13). Os pacientes fisicamente inativos apresentavam maior peso corporal, maior GC e menor MMC em % de peso corporal quando comparados com indivíduos ativos. Os pacientes obesos (com base no IMC) tiveram pior nível de AFVD do que paciente subnutridos, eutróficos e sobrepeso. Piores níveis de AFVD também foram observados em pacientes com obesidade mórbida com base na %GC apesar da modesta correlação entre GC e TAF>moderada no grupo completo. **Conclusões:** Pacientes com DPOC fisicamente ativos na vida diária possuem proporcionalmente mais MMC e menos GC do que pacientes fisicamente inativos. Inatividade física mais acentuada é observada em pacientes obesos, apesar da composição corporal em geral não poder ser considerada um fator fortemente correlacionado aos níveis de AFVD de pacientes com DPOC.

Palavras-chave: Doença pulmonar obstrutiva crônica. Obesidade. Composição corporal. Exercício. Atividade física.

MONTEIRO, Fabiane. **Obesity and physical activity in daily life in patients with COPD**. 2011. 43 p. Trabalho de Conclusão de Curso do Programa de Pós-Graduação em Ciências da Reabilitação (Programa Associado entre Universidade Estadual de Londrina [UEL] e Universidade Norte do Paraná [UNOPAR]) – Universidade Estadual de Londrina, Londrina, 2011.

ABSTRACT

Purpose: To study differences in body composition in physically active and inactive patients with chronic obstructive pulmonary disease (COPD); and to study the relationship of obesity with physical activity in daily life (PADL) in this population.

Methods: Body composition (bioelectrical impedance analysis), and level of PADL (activity monitors DynaPort and SenseWear) were evaluated in 74 patients with COPD (45 men, 65±9years, FEV₁=40±15%pred, IMC=27±6kg.m⁻²). Patients were separated in 2 groups: physically active (> 30 minutes/day of time spent in physical activity [TPA] of at least moderate intensity, or TPA>moderate) and inactive (did not achieve these recommendations). The sample was also classified according to the body mass index (BMI) in underweight, normal weight, overweight and obese; and according to their percentage of fat mass (%FM) in moderate obesity, high obesity and morbid obesity. **Results:** In the whole group, TPA>moderate correlated weakly with BMI (r=-0.28; p=0.02) and FM (r=-0.30; p=0.001) but not with FFM (r=-0.18; p=0.13). Physically inactive patients had higher body weight and higher FM and lower fat free mass (FFM) in % of body weight than active patients. Obese patients (according to BMI) had worse PADL level than underweight and normal weight patients. Poorer PADL also occurred in morbidly obese patients classified according to %FM, despite the weak correlation between FM and TPA>moderate in the whole group. **Conclusion:** Physically active patients with COPD have proportionally more FFM and less FM than inactive patients. More pronounced physical inactivity occurs in obese patients, although body composition in general does not qualify as an important correlate factor of the level of PADL in patients with COPD.

Key words: Chronic obstructive. Pulmonary disease. Obesity. Body composition. Exercise. Physical activity.

LISTA DE TABELAS

Tabela 1 – Resultados gerais do grupo de pacientes com doença pulmonar obstrutiva crônica (DPOC) e dos grupos de pacientes fisicamente ativos e inativos baseado nas recomendações do American College of Sports Medicine (ACSM).....	29
Tabela 2 – Comparação de pacientes com DPOC classificados de acordo com índice de massa corpórea (IMC).....	30
Tabela 3 – Comparação de pacientes com DPOC classificados de acordo com o percentual de gordura corporal (GC%).....	30

LISTA DE ABREVIATURAS E SIGLAS

6MWT	6-Minute Walking Test
%FM	Percentage of Fat Mass
%GC	Percentual de Gordura Corporal
ACSM	American College of Sports Medicine
AFVD	Atividade Física na Vida Diária
BIA	Bioelectrical Impedance Analysis
BMI	Body Mass Index
COPD	Chronic Obstructive Pulmonary Disease
DAM	DynaPort Activity Monitor
DPOC	Doença Pulmonar Obstrutiva Crônica
FFM	Fat Free Mass
FFMI	Fat Free Mass Index
FM	Fat Mass
GC	Gordura Corpórea
GOLD	Global Initiative for Chronic Obstructive Lung Disease
HO	High Obesity
IMC	Índice de Massa corpórea
LFIP	Laboratório de Pesquisa em Fisioterapia Pulmonar
MbO	Morbid Obesity
MET	Metabolic Equivalent
MMC	Massa Magra Corpórea
MO	Moderate Obesity
NW	Normal Weight
OB	Obese
OMS	Organização Mundial da Saúde
OW	Overweight
PADL	Physical Activity in Daily Life
SAB	SenseWear Armband Activity Monitor
TAF>moderada	Tempo Gasto Por Dia Em Atividade Física de intensidade ao menos moderada
TPA>moderate	Time Spent in Physical Activity Of At Least Moderate Intensity
TW	Time Spent Walking per day

UEL	Universidade Estadual de Londrina
UW	Underweight
VEF₁/CVF	Razão entre Volume Expiratório Forçado no Primeiro Secundo (VEF ₁) e Capacidade Vital Forçada (CVF)

SUMÁRIO

1	INTRODUÇÃO	11
2	REVISÃO DE LITERATURA – CONTEXTUALIZAÇÃO	13
2.1	DOENÇA PULMONAR OBSTRUTIVA CRÔNICA.....	13
2.2	OBESIDADE	14
2.3	NÍVEL DE ATIVIDADE FÍSICA	16
2.4	RELAÇÃO ENTRE DPOC, OBESIDADE E NÍVEL DE ATIVIDADE FÍSICA	18
3	ARTIGO	20
4	CONCLUSÕES GERAIS	31
	REFERÊNCIAS	32
	ANEXOS	37
	ANEXO A – Normas de formatação do periódico LUNG	38

1 INTRODUÇÃO

A doença pulmonar obstrutiva crônica (DPOC) é caracterizada por comprometimentos pulmonares e extra-pulmonares. Além da limitação ao fluxo aéreo e dispnéia, a doença cursa com efeitos sistêmicos como redução da capacidade de exercício, força muscular e qualidade de vida (1). Alterações na composição corporal como redução de índice de massa corpórea (IMC) e de massa magra corpórea (MMC) são comumente observados, além de serem preditores independentes de mortalidade na doença (2-4). No entanto, com o crescimento dos índices mundiais de obesidade, inclusive em pacientes com DPOC (5;6), maior atenção tem sido dada não somente a pacientes com DPOC caracterizados por caquexia mas também aos pacientes considerados obesos.

Sabe-se que pacientes obesos apresentam alterações respiratórias importantes como aumento na demanda e no trabalho ventilatório, diminuição da complacência torácica e conseqüentemente maior dispnéia, tudo isso devido ao aumento de gordura acumulada em região torácica e abdominal (7). Essas alterações respiratórias associadas a problemas ortopédicos comuns da obesidade como dores articulares, lombalgias e edema, fazem com que a atividade física seja dificultada nesta população. Estudos mostram que pacientes com DPOC obesos (com base no IMC) possuem pior capacidade funcional de exercício quando comparados a pacientes com IMC normal (5;8).

As diretrizes de saúde pública a respeito da atividade física publicadas pelo American College of Sports Medicine (ACSM) recomendam que um mínimo de 30 minutos de atividade física de intensidade moderada é necessária para a manutenção ou desenvolvimento da capacidade física (9). Estudos anteriores mostraram que o nível de atividade física na vida diária (AFVD) é reduzido em pacientes com DPOC e que essa redução está associada a maiores índices de mortalidade (10).

Um estudo realizado por Park e cols (11) com mulheres saudáveis mostrou que aquelas que apresentavam maior percentual de gordura corporal apresentavam menor número de passo por dia e menos tempo em atividade física de intensidade moderada e vigorosa. O estudo discute ainda que a relação entre obesidade e atividade física diária deve ser discutida usando não apenas o IMC mas também o índice de massa de gordura corporal (GC) em kilos ou o percentual de

gordura corporal (%GC), já que o IMC não expressa necessariamente a quantidade de GC. No entanto, ainda não se sabe se existe relação entre obesidade e nível de AFVD em pacientes com DPOC e se pacientes com maiores níveis de obesidade (com base no %GC) possuem piores níveis de AFVD. Sendo assim, os objetivos deste estudo foram: I) determinar se existem diferenças na composição corporal entre os pacientes com DPOC fisicamente ativos e inativos (de acordo com as recomendações da ACSM); II) verificar se os pacientes obesos (classificados pelo IMC) têm menor nível de AFVD do que pacientes subnutridos, eutróficos e sobrepeso; III) investigar se existem diferenças na capacidade funcional de exercício e nível de AFVD entre pacientes com diferentes níveis de obesidade determinada de acordo com a %GC.

2 REVISÃO DE LITERATURA – CONTEXTUALIZAÇÃO

2.1 DOENÇA PULMONAR OBSTRUTIVA CRÔNICA

Segundo o documento da **Global Initiative for Chronic Obstructive Lung Disease (GOLD)**, a Doença Pulmonar Obstrutiva Crônica (DPOC) é definida como uma doença prevenível e tratável com alguns efeitos extrapulmonares significantes que podem contribuir para a gravidade individualmente. A alteração pulmonar característica da doença é a limitação ao fluxo aéreo, que não é totalmente reversível. Esta limitação é geralmente progressiva e associada à resposta inflamatória anormal dos pulmões a partículas nocivas ou gases (1).

A Organização Mundial de Saúde (OMS) estima que a DPOC atinja 210 milhões de pessoas em todo o mundo, sendo dentre essas cerca de 64 milhões casos sintomáticos. A prevalência da doença é muito variável nas diferentes regiões do mundo, sendo as Américas responsáveis por 13,2 milhões dos casos sintomáticos (12). A DPOC está entre as maiores causas de morbidade e mortalidade em todo o mundo (13), e em geral está associada ao consumo do tabaco (1), sendo este responsável por cerca de 90% dos casos da doença. Segundo estimativa do Global Burden of Disease Study, devido à expansão da epidemia do tabagismo e às mudanças demográficas na maioria dos países (e.g., aumento da expectativa de vida), em 2020 a DPOC será a terceira principal causa de morte do mundo (14) e a quarta mais importante doença que leva a incapacidade física (1). O cigarro é o principal fator de risco para desenvolvimento da DPOC; no entanto, outros fatores como poluição ambiental, produtos químicos e tóxicos e até mesmo componentes genéticos podem estar associados ao desenvolvimento da doença. Devido ao seu caráter progressivo e incapacitante, acarreta um considerável impacto econômico e social, pela redução na produtividade, mortes prematuras, comprometimento do orçamento familiar, aposentadorias precoces e alto custo com o tratamento e com as internações, que são freqüentes (15). Quanto à morbidade nesses pacientes, apesar da literatura não apresentar dados claros e objetivos, estima-se que o aumento da idade e o sexo masculino estejam mais relacionados a maiores níveis de morbidade (16). No entanto, a tendência de prevalência para os homens, especialmente os mais jovens, é de estabilidade e até mesmo decréscimo, enquanto que para as mulheres é de crescimento (17). Dados

de prevalência e morbidade podem ainda ser subestimados já que os pacientes são diagnosticados somente em fases mais avançadas da doença, quando os sinais clínicos começam a aparecer.

Em geral, a DPOC costuma ser diagnosticada por volta da quinta ou sexta década de vida (18) e o diagnóstico é feito com base em variáveis espirométricas. A razão volume expiratório forçado no primeiro segundo (VEF_1) / capacidade vital forçada (CVF), ou $VEF_1/CVF < 70\%$ do predito em combinação com $VEF_1 < 80\%$ do predito (pós-broncodilatador) em indivíduos que apresentam tosse produtiva, dispnéia ou exposição a fatores de risco confirmam o diagnóstico(1). No entanto, a avaliação da DPOC exclusivamente pela alteração pulmonar tem sido vista pela literatura como insuficiente, já que esta é reconhecidamente uma doença que cursa com alterações extrapulmonares significativas (19). Além da dispnéia, tosse, sibilância, produção de secreção e infecções respiratórias de repetição, consequências sistêmicas tais como descondicionamento, fraqueza muscular, inatividade física na vida diária, perda de peso, e desnutrição são frequentemente observadas (20;21). Problemas emocionais como depressão, ansiedade e isolamento social também são comumente encontrados em pacientes com DPOC (22). Os preditores de mortalidade na DPOC são maior idade e histórico tabágico, hipoxemia dispnéia e hipersecreção crônica mais acentuadas, maior redução na capacidade de exercício, atividade física na vida diária, massa e força muscular, baixo índice de massa corpórea e perda de peso excessiva (23).

2.2 OBESIDADE

A Organização Mundial de Saúde define a obesidade como o excesso de gordura corporal que gera implicações negativas para a saúde (24). A obesidade é uma condição complexa de dimensões sociais, biológicas e psicossociais consideráveis, podendo eventualmente afetar qualquer pessoa de qualquer idade ou grupo socioeconômico, em qualquer parte do mundo. É hoje considerada um fenômeno global que aumenta consideravelmente a morbidade e diminui expectativa de vida (7), sendo seu impacto mais pronunciado na morbidade do que na mortalidade (25).

Segundo a OMS, a prevalência da obesidade mais que dobrou desde 1980. Em 2008, 1,5 bilhões de adultos acima de 20 anos foram considerados

sobrepeso, e dentre estes, mais de 200 milhões de homens e aproximadamente 300 milhões de mulheres foram considerados obesos (24). Dados mais recentes da OMS, mostram que 11,1% dos brasileiros adultos são considerados obesos ($IMC \geq 30 \text{ Kg.m}^{-2}$) (26).

A obesidade é causada por um desequilíbrio entre ingestão e consumo calórico. Está relacionada a fatores genéticos, que são considerados não modificáveis, e fatores modificáveis como consumo calórico excessivo, sedentarismo e fatores psicológicos. Algumas doenças como hipotireoidismo, Síndrome de Cushing e até mesmo medicações como esteróides e antidepressivos podem predispor o indivíduo a desenvolver obesidade.

A obesidade está fortemente relacionada a diversas comorbidades como hipertensão, doença arterial coronariana, câncer, apnéia obstrutiva do sono e osteoartrose, contribuindo diretamente para maior gravidade destas (27-29). É considerada ainda o maior fator de risco para doenças crônicas incluindo doenças cardiovasculares, cerebrovasculares e diabetes mellitus (30). O sistema respiratório também é bastante prejudicado pela obesidade. Resumidamente, a obesidade pode reduzir um terço da complacência torácica devido ao acúmulo de gordura torácica e abdominal (31), gerando aumento do trabalho respiratório e conseqüentemente ocasional dispnéia (32).

A forma mais comum de se graduar a obesidade é por meio da utilização do índice de massa corpórea (IMC). Este índice é internacionalmente conhecido (33) e amplamente utilizado em estudos epidemiológicos tendo sido incorporado à prática clínica devido à sua simplicidade e praticidade. No entanto, o IMC é uma relação simples entre peso corporal e altura, e portanto não é capaz de distinguir a gordura corporal da massa muscular (34). Sabendo-se que a massa muscular não causa prejuízo algum à saúde do indivíduo e que somente a gordura corporal está relacionada aos efeitos deletérios da obesidade (35), a medida exclusiva da gordura corporal parece ser melhor preditiva de obesidade do que o IMC, que leva em consideração apenas o peso corporal como um todo. Além disso, a literatura mostra que o IMC é considerado impreciso para detectar o percentual de gordura especialmente em populações com diferentes idades e tipos corporais (36). No entanto, apesar de todas essas limitações, o IMC continua sendo a ferramenta mais utilizada para classificar a obesidade devido à sua praticidade.

2.3 NÍVEL DE ATIVIDADE FÍSICA NA VIDA DIÁRIA

A atividade física é definida como qualquer movimento corporal produzido pelos músculos esqueléticos que resulta em gasto energético maior do que os níveis de repouso (37). Segundo a Organização Mundial de Saúde, a prática de atividade física regular reduz o risco de mortes prematuras, doenças do coração, acidente vascular cerebral, diabetes tipo II e câncer de cólon e de mama (38). Atua também na prevenção ou redução da hipertensão arterial, previne o ganho de peso (diminuindo o risco de obesidade), auxilia na prevenção ou redução da osteoporose, promove bem-estar, reduz o estresse, a ansiedade e a depressão. Especialmente em crianças e jovens, a atividade física interage positivamente com as estratégias para adoção de uma dieta saudável, desestimula o uso do tabaco, álcool e drogas, reduz a violência e promove a integração social (38). A atividade física regular está associada ainda à redução nos índices de morbidade e mortalidade, menor limitação funcional assim como menor número de quedas em idosos (39). Um estudo realizado pelo European Community Respiratory Health Survey II (ECRHS-II) mostrou que a inatividade física é um importante fator de risco para desenvolvimento de hiperresponsividade brônquica na população adulta em geral (40).

O sedentarismo é hoje considerado um problema mundial de saúde e está relacionado a altos custos com tratamentos, inclusive e principalmente hospitalares. Segundo dados do Centers for Disease Control and Prevention, só no ano de 2000 nos Estados Unidos o sedentarismo foi responsável pelo gasto de 76 bilhões de dólares com custos médicos, mostrando assim que seu combate merece prioridade na agenda de saúde pública (41). Segundo dados da OMS (42), a inatividade física é o quarto principal fator de risco para mortalidade. Aproximadamente 3,2 milhões de mortes por ano são atribuídas aos baixos níveis de atividade física. O rápido crescimento das doenças crônicas associadas à inatividade física vem sendo registrado tanto nos países desenvolvidos como nos países em desenvolvimento, sendo que a maioria dos adultos nos países desenvolvidos é considerada inativa (42). A atividade física regular tem sido considerada como um dos principais componentes na prevenção do crescimento global de doenças crônicas.

As diretrizes de saúde pública a respeito da atividade física publicadas pelo American College of Sports Medicine (ACSM) (9) preconizam que

sessões de trinta minutos de atividades físicas de intensidade moderada por dia, na maior parte dos dias da semana, desenvolvidas continuamente ou mesmo em períodos cumulativos de 10 minutos, já são suficientes para a manutenção ou para o desenvolvimento da aptidão física, independentemente da idade. Indivíduos que não atingem estes padrões mínimos são considerados insuficientemente ativos e apresentam um maior risco de morbi-mortalidade.

Um estudo feito nos EUA em 2005 mostrou que menos da metade dos adultos (41,9%) atingiram as recomendação de atividade física recomendadas pelo ACSM e que os homens cumpriam mais as recomendações em comparação com as mulheres (50,7% e 47,9%, respectivamente). Além disso, o estudo mostra que os jovens foram mais ativos que os mais velhos e que pessoas com nível de escolaridade maior são os que mais atingem as recomendações (41). Pessoas que não atingem as recomendações mínimas (ACSM) possuem 20 a 30% mais chance de mortalidade. Em 2008, mundialmente, 31% dos adultos com idade superior a 15 anos foram considerados inativos (28% dos homens e 34% das mulheres), sendo a região das Américas e Leste do Mediterrâneo as com maior prevalência de inatividade física. Em todos os países, os homens foram mais ativos que as mulheres. No Brasil, 49,4 % dos indivíduos foram considerados insuficientemente ativos (43).

Existem diversas formas de se quantificar o nível de atividade física na vida diária (AFVD). As principais formas são os questionários de auto-relato, os sensores de movimento como acelerômetros e pedômetros e métodos que mensuram diretamente o gasto energético como a técnica *doubly labeled water* (44;45). Os questionários de auto-relato são bastante suscetíveis a viés, já que dependem da memória do avaliado. A literatura mostra que os questionários se correlacionam fracamente com a quantificação objetiva da AFVD (46;47) e não fornecem dados precisos de gasto energético (45). Pedômetros são sensores de movimento portáteis, baratos e de fácil manuseio; no entanto, fornecem somente informações básica como contagem de número de passos e não são eficazes na detecção de passos em intensidades baixas de movimento, fazendo com que seu uso seja questionável para determinadas populações (48), como pacientes com DPOC. Já os acelerômetros multiaxiais como o DynaPort (McRoberts, Holanda) e SenseWear Armband (BodyMedia, Estados Unidos da América), apesar de apresentarem um custo mais elevado, fornecem informações detalhadas sobre o

padrão, tempo e intensidade da atividade física (47;49). Além disso, os sensores de movimento multiaxiais são mais úteis para avaliação de populações mais inativas, já que são mais sensíveis na avaliação de movimentos de baixa intensidade (44).

2.4 RELAÇÃO ENTRE DPOC, OBESIDADE E NÍVEL DE ATIVIDADE FÍSICA NA VIDA DIÁRIA

Tanto a obesidade como a inatividade física são fatores de risco modificáveis para diversas doenças crônicas e estão associados ao alto grau de inflamação sistêmica, o que pode potencializar efeitos negativos sobre a saúde e bem estar dos portadores destas doenças (50). Alterações na composição corporal e no nível de AFVD são preditores independentes de morbi-mortalidade em pacientes com DPOC (3;51-53). Assim como na população geral, estudos têm demonstrado que a obesidade vem crescendo exponencialmente em pacientes com DPOC (5;6). A obesidade está relacionada a diversas alterações respiratórias como aumento da demanda e do trabalho ventilatório, redução na complacência torácica e conseqüentemente maior grau de dispnéia (7). Um estudo publicado por Naimark e Cherniack mostrou que a complacência torácica em indivíduos obesos é reduzida a dois terços de um indivíduo normal, e tal fato é decorrente principalmente do acúmulo de gordura torácica e abdominal (31). A fraqueza muscular respiratória também está comumente presente na obesidade e tem sido atribuída à ineficiência muscular decorrente da redução da complacência torácica, do volume pulmonar ou ambos (54;55). Alterações respiratórias associadas a problemas ortopédicos, como por exemplo dor articular, que está comumente presente na obesidade, faz com estes paciente reduzam sua capacidade de exercício. Ramachandram e cols (5), realizaram um estudo com pacientes com DPOC obesos e demonstraram que apesar destes pacientes serem encaminhadas para reabilitação em estágios mais iniciais da doença (de acordo com o GOLD), estes apresentam pior capacidade funcional de exercício, maior grau de comprometimento funcional e maiores níveis de fadiga. No entanto, este mesmo estudo mostra que apesar de apresentarem piores valores basais, pacientes com DPOC considerados obesos se beneficiam igualmente do programa de reabilitação pulmonar. Seres e cols (56) atribuem essa redução da capacidade de exercício à necessidade de maior gasto energético para mover a massa corporal aumentada. Sabe-se ainda, que pacientes com DPOC apresentam menores níveis de AFVD quando comparados a indivíduos saudáveis

pareados pela idade (21) e que na população geral o baixo nível de atividade física está entre os principais fatores predisponentes à obesidade. Apesar de diversos estudos terem mostrado que pacientes com DPOC obesos possuem pior capacidade de exercício, ainda não se sabe se esta mesma relação se aplica ao nível de AFVD.

3 ARTIGO:

OBSESITY AND PHYSICAL ACTIVITY IN DAILY LIFE IN PATIENTS WITH COPD

(Em revisão no periódico *LUNG*)

Fabiane Monteiro¹; Carlos Augusto Camillo^{1,2}; Renato Vitorasso¹; Thaís Sant'Anna¹; Nídia A. Hernandez¹; Vanessa S. Probst^{1,3}; Fábio Pitta^{1*}.

***Corresponding author:**

Dr. Fabio Pitta

Departamento de Fisioterapia – CCS, Hospital Universitário de Londrina

Rua Robert Koch, 60 – Vila Operaria, 86038-350 – Londrina, Paraná, Brazil.

Phone.: +55 43 3371 2288; Fax: +55 43 3371 2459.

E-mail: fabiopitta@uol.com.br

Keywords: Chronic obstructive. Pulmonary disease. Obesity. Body mass index. Body composition. Exercise. Physical activity.

Abstract: Purpose: To study differences in body composition in physically active and inactive patients with COPD; and to study the relationship of obesity with physical activity in daily life (PADL) in this population. **Methods:** Body composition (bioelectrical impedance analysis), and level of PADL (activity monitors DynaPort and SenseWear) were evaluated in 74 patients with COPD (45 men, 65±9years, FEV₁=40±15%pred, IMC=27±6kg.m⁻²). Patients were separated in 2 groups: physically active (> 30 minutes/day of time spent in physical activity [TPA] of at least moderate intensity, or TPA>moderate) and inactive (did not achieve these recommendations). The sample was also classified according to the body mass index (BMI) in underweight, normal weight, overweight and obese; and according to their percentage of fat mass (%FM) in moderate obesity, high obesity and morbid obesity. **Results:** In the whole group, TPA>moderate correlated weakly with BMI (r=-0.28; p=0.02) and FM (r=-0.30; p=0.001) but not with FFM (r=-0.18; p=0.13). Physically inactive patients had higher body weight and FM and lower fat free mass (FFM) in % of body weight than active patients. Obese patients (according to BMI) had worse PADL level than underweight and normal weight patients. Poorer PADL occurred in morbidly obese patients, despite the weak correlation between FM and TPA>moderate in the whole group. **Conclusion:** Physically active patients with COPD have proportionally more FFM and less FM than inactive patients. More pronounced physical inactivity occurs in obese patients, although body composition does not qualify as an important correlate factor of the level of PADL in patients with COPD.

¹ Laboratório de Pesquisa em Fisioterapia Pulmonar (LFIP), Departamento de Fisioterapia, Universidade Estadual de Londrina (UEL), Londrina, Paraná, Brazil

² Programa de Mestrado em Fisioterapia, Univ Estadual Paulista (UNESP), Presidente Prudente, São Paulo, Brazil

³ Centro de Pesquisa em Ciências da Saúde, Centro de Ciências Biológicas e da Saúde, Universidade Norte do Paraná (UNOPAR), Londrina, Paraná, Brazil

Introduction

Chronic Obstructive Pulmonary Disease (COPD) is characterized by airflow limitation, dyspnea and reduction in exercise capacity, muscle strength and quality of life[1]. Patients with COPD frequently also present alterations in their body composition such as weight loss and fat-free mass depletion, and it is known that reduced body mass index (BMI) and fat free mass index (FFMI) are independent predictors of mortality in these patients[2-5]. On the other hand, recent studies indicate that obesity has been increasingly common in patients with COPD[6-8]. The body mass index (BMI) is the most used tool to identify obesity in any population. It is an internationally known index, used in clinical practice due to its low cost and simplicity. However, the BMI does not distinguish fat mass (FM) from fat free mass (FFM) since it is just a simple ratio between body weight and height squared. Therefore, since obesity is usually considered as the excess of FM, the quantification of FM is another option for the identification of obesity, The major respiratory complications of obesity include an increased ventilatory demand and work of breathing, respiratory muscle inefficiency and reduced respiratory compliance[9], all contributing to an increase in dyspnea. These respiratory changes associated with muscle loss, joint pain and skin friction are important determinants of decreased exercise capacity and disability in obese patients with COPD[9]. Despite previous studies suggesting that obese patients with COPD have lower functional exercise capacity[6], there are no studies that show the relationship between obesity and level of physical activity in daily life (PADL) in these patients. PADL is normally reduced in patients with COPD[10;11] and as described for the alterations in body composition, reduction in PADL is a modifiable morbi-mortality risk factor[12;13].

Public health guidelines regarding physical activity published by the American College of Sports Medicine (ACSM) recommend that a minimum of 30 minutes of physical activity of moderate intensity (e.g., walking) is necessary for the maintenance or development of physical fitness[14]. It is yet unknown whether patients with COPD who are physically active according to these recommendations have different body composition in comparison to the physically inactive ones. Therefore, the present study had three aims: to determine if there are differences in body composition between active and inactive patients with COPD (according to the ACSM recommendations); to verify whether obese patients (according to BMI) have lower level of PADL than underweight, normal weight and overweight patients; and to investigate if there are differences in functional exercise capacity and PADL between levels of obesity according to the percentage of fat mass.

Methods

Sample, inclusion and exclusion criteria

Ninety patients with COPD (55 men, 66±9 years, forced expiratory volume in the first second (FEV₁) = 42±16%pred, BMI=27±8kg.m⁻²) were initially included in this cross-sectional retrospective study. Patients were recruited during the procedure of admission to a Pulmonary Rehabilitation Program of a University Hospital (State University of Londrina, Brazil). Since 16 subjects did not have complete data of the PADL, they were not included at the analysis. Diagnosis of COPD was based on the Global Initiative for

Chronic Obstructive Lung Disease spirometry criteria[1], i.e., post-bronchodilator (Aerolin®) ratio between FEV₁ and forced vital capacity (FVC) < 0.70. As inclusion criteria, patients should be in stable condition (no exacerbations in the last three months that required a change in medication or hospital admission); should not have participated in any exercise training program in the last year before admission to the study; and could not present other pathological conditions which could impair their performance, such as cerebro-vascular, orthopedic or rheumatic disease. Exclusion criteria were the occurrence of severe acute exacerbation requiring hospitalization during the assessment period; and the inability to understand or cooperate with the assessment methods. The ethics committee of Londrina University Hospital approved the research protocol and a written consent form was obtained from each patient.

Assessments

Pulmonary function (spirometry) was performed using a Pony Graphics® spirometer (Cosmed, Rome, Italy) according to international recommendations[15]. Reference values used were those of Pereira et al[16].

Functional exercise capacity was evaluated by the 6-minute walking test (6MWT), in accordance with standards of the American Thoracic Society[17]. Reference values were those described by Troosters et al[18].

Body composition analysis was performed by bioelectrical impedance (Biodynamics 310, Biodynamics, BR) (BIA), using the technique described by Lukaski et al[19]. Reference values used were those described specifically for patients with chronic lung disease by Kyle et al[20]. Body mass index (BMI) was calculated by dividing weight by height squared, and fat free mass index (FFMI) by dividing the weight of fat free mass (obtained by BIA) by height squared.

Assessment of physical activity in daily life was performed simultaneously by two activity monitors: the accelerometer-based DynaPort Activity Monitor, or DAM (McRoberts BV; The Hague, the Netherlands) and the multisensor SenseWear Armband, or SAB (BodyMedia Inc., Pittsburgh, PA, USA). The DAM consists of a small, lightweight box enclosed in a belt worn around the waist and a leg sensor (total weight, 375 g). The DAM was shown to be as accurate as video recordings (criterion method) in order to assess time spent in different activities in patients with COPD, such as the time spent walking/day (TW)[21]. Technical specifications about the DAM can be found elsewhere[21]. The SAB is a lightweight (80g) activity monitor, worn on the triceps brachial bulk of the right arm. The device estimates energy expenditure based on a biaxial accelerometer and physiologic sensors that detect galvanic skin response, heat flux and skin temperature in a manufacturer algorithm. It also provides the duration of activities performed above a determined level of intensity (e.g. time spent per day in activities of at least moderate intensity, or TPA>moderate). The SAB was already validated for the estimative of energy expenditure in patients with COPD[22-25]. Main outcomes of the PADL assessment were TW (from the DynaPort) and TPA>moderate (from the SenseWear), since these outcomes are complementary and provide together an overall view of the patient's PADL level, both regarding time and intensity. Assessments with both activity monitors were performed during 2 consecutive weekdays, for 12 hours per day (from wake-up time to 12h after awaking). The mean of the two assessment days was used for statistical analysis, and the minimal number of days needed to obtain a reliable assessment of daily physical

activity was determined in a previous study[10]. For analysis, patients were first divided into two groups, according to the minimum recommendation of daily physical activity determined by the ACSM[14]: active (performed more than 30 minutes/day of TPA>moderate) and inactive (did not achieve these recommendation). Threshold of metabolic equivalents (METs) to characterize an activity as being of at least moderate intensity was also suggested by the ACSM: above 4.8 METs for subjects aged 28-39 years; above 4.0 METs for 40-60 years; above 3.2 METs for 65-79 years and above 2.0 METs for subjects over 80 years old[14]. Secondly, the same group was divided according to the BMI classification suggested for patients with COPD by the ATS-ERS Statement of Pulmonary Rehabilitation[26] into underweight [UW] ($BMI < 21 \text{ Kg.m}^{-2}$), normal weight [NW] ($21 \leq BMI < 25 \text{ Kg.m}^{-2}$), overweight [OW] ($25 \leq BMI < 30 \text{ Kg.m}^{-2}$) and obese [OB] ($BMI > 30 \text{ Kg.m}^{-2}$). Finally, the group was separated according to the percentage of fat mass (%FM) into non-obese (women [W]: $< 25\%FM$; men [M]: $< 15\%FM$); mild obesity (W: $25 \leq \%FM < 30$; M: $15 \leq \%FM < 20$); moderate obesity [MO] (W: $30 \leq \%FM < 35$; M: $20 \leq \%FM < 25$), high obesity [HO] (W: $35 \leq \%FM < 40$; M: $25 \leq \%FM < 30$) and morbid obesity [MbO] (W: $\%FM \geq 40$; M: $\%FM \geq 30$)[27].

Statistical Analysis

Statistical analysis was performed using the SPSS statistical package software version 17.0 (SPSS inc., Chicago IL) and GraphPad Prism 3.0 (GraphPad Software, San Diego, CA, USA) programs. Kolmogorov-Smirnov test was used to check for normal distribution. Comparisons between groups were performed using Student t test or Mann-Whitney test; and one-way ANOVA or Kruskal-Wallis, followed by a post-hoc test (Bonferroni or Dunns, respectively). Correlations were studied using the Pearson or Spearman coefficient, according to the normality in data distribution. To verify the similarity in gender between the groups, the chi-square test was used. A significance level of $p \leq 0.05$ was adopted.

Results

Sixteen subjects were excluded since complete data of the PADL assessment could not be obtained due to technical problems or misuse of the device by the patient. Therefore, for the statistical analysis, the sample was composed by 74 subjects.

According to the ACSM guidelines, 70% of the 74 patients included in the study were classified as physically inactive (52 *versus* 22 physically active). Results of the whole group and differences in body composition between physically active and inactive subgroups are shown in Table 1. Physically active and inactive patients with COPD had important differences in their body composition. Inactive patients had higher body weight ($p=0.002$), BMI ($p=0.001$) and FM ($p=0.0005$). Due to the higher body weight, the inactive group had higher absolute values of FFM ($p=0.05$). However, when FFM was corrected for body weight ($\%FFM = [FFM \text{ in Kg} \times 100] / \text{body weight in Kg}$), the inactive group had significantly lower values ($p=0.03$).

In the whole group, TPA>moderate correlated weakly and negatively with weight ($r=-0.29$; $p=0.02$), BMI ($r=-0.28$; $p=0.02$) and FM ($r=-0.30$; $p=0.001$), but not with FFM ($r=-0.18$; $p=0.13$). Correlation between BMI and %FM was $r=0.47$; $p<0.0001$. The 6MWT was significantly lower in inactive patients, whereas the active and inactive subgroups did not differ in their airway obstruction (table 1).

When the whole group was divided according to BMI, 19% of the patients were UW; 23% were NW; 30% were OW; and 28% were OB. Table 2 shows that, when comparing the level of PADL between the four groups, OB had significantly lower TW than UW and NW and significantly lower TPA>moderate than UW ($p<0.05$ for all). No differences were found between OW with the other three groups (UW, NW and OB). Despite the differences in PADL between OB *versus* UW and NW, the groups did not differ significantly in the 6MWT, despite an average difference of 54 meters between OB and UW and 41 meters between OB and NW.

As for the classification according to the %FM, no subject in the present sample could be classified as non-obese. Three patients had mild obesity and were therefore excluded from the subsequent analysis. Table 3 offers a detailed overview of the differences between MO, HO and MbO in patients with COPD. Concerning PADL, significant differences were found in TW (MbO *vs* MO and *vs* HO) and TPA>moderate (MO *vs* MbO and *vs* HO). Poorer 6MWT occurred in morbidly obese patients (table 3).

Two retrospective (*post-hoc*) power calculations were performed based on the observed differences in the two main outcomes of physical activity in daily life (TPA>moderate and TW) between the obese ($n=21$) and underweight groups ($n=14$) (table 2). Taking into consideration an alpha of 0.05, a difference of 31 minutes in TPA>moderate and 27 minutes in TW between the groups and a standard deviation of 28 minutes (for both outcomes), the study had a power of 87% to detect a significant difference in TPA>moderate and 78% to detect a significant difference in TW.

Discussion

This study showed that despite physically active and inactive patients with COPD have similar age, airway obstruction and gender proportion, inactive patients have less FFM and more FM (in % of body weight) than their active counterparts. In addition, time spent in activities of at least moderate intensity (as classified by the ACSM) has weak correlation with BMI and absolute values of FM but no correlation with FFM. Almost 30% of the patients were obese according to the classification based on the BMI. On the other hand, when using %FM to classify obesity, 100% of patients were classified as obese, either mildly, moderately, highly or morbidly. Obese patients presented worse level of daily physical activity than underweight and normal weight patients. Moreover, patients with higher degrees of obesity (according to %FM) had more impaired PADL and worse functional exercise capacity.

It is widely known that patients with COPD have important alterations in body composition and this is an independent predictor of mortality[4]. We now observed that inactive patients with COPD had higher body weight and BMI, and this was mainly caused by higher weight of fat mass. The inactive subgroup had higher absolute values of FFM, but this was due to the fact that these subjects had higher body weight, since FFM in % of body weight was significantly lower in this group. Moreover, although weakly, fat mass was better correlated with the level of PADL than did fat free mass. These results suggest that, in COPD, obesity may be somewhat better linked to physical inactivity than loss of FFM, although none of the two factors appear to be a strong correlate of the level of PADL in this population. This corroborates previous findings by Watz et al[28], who studied the determinants of objectively measured physical activity in patients with COPD. These authors also concluded that BMI was not a determinant factor for PADL level. Other studies also did not find BMI to be a strong correlate of the level of physical activity in daily life in patients with COPD[10;29;30].

Obesity is known as an important issue in COPD and BMI is yet the most used variable to identify obesity in these patients. Literature reported obesity prevalence (according to BMI) in COPD as ranging between 18%[31] to 54%[32]. In the present study 28% of the patients were obese according to their BMI. Our results showed that obese patients ($BMI \geq 30 \text{Kg.m}^{-2}$) are less active in comparison to underweight and normal-weight patients with COPD, but not in comparison to overweight patients. Similar results were found in a study with older healthy adults, in which obesity but not overweight was associated with lower levels of physical activity[33].

Although differences in the 6MWT between the groups classified according to BMI did not reach statistical significance, our data showed that obese patients walked on average 54 meters less than underweight patients and 41 meters less than normal weight patients, which are both well above the two values of minimal clinically important difference recently suggested for the 6MWT in the literature (25 meters and 35 meters)[34;35]. Therefore, despite not reaching statistical significance, a worse functional exercise capacity in obese patients appears to be clinically relevant. The previous literature has shown that higher BMI is associated with lower functional exercise capacity[8;9;36] and this is likely the result of the increase in energy expenditure associated with weight bearing exercise[37].

The BMI is the most used outcome to classify obesity, although does not provide clear information about the distribution of FM and FFM. Patients with COPD commonly develop cachexia which is caused by different factors such as increased metabolic rate, hypoxaemia, sympathetic upregulation, inactivity, oxidative stress, inflammation and anabolic hormone insufficiency[38]. This reduction in percentage of FFM leads to higher proportion of fat mass, even though the patient's weight is not necessarily high. This explains the fact that 100% of the present sample was classified as obese based on the % of FM, being 42% of them classified as morbidly obese. A study by Park et al.[33], which verified the relation of body composition with PADL in adult women, suggested that the relationship between obesity and daily physical activity should be discussed using not only BMI but also fat mass index or %FM since these two variables were related to the physical activity level, what was corroborated by the present results. Obesity is defined as excessive accumulation of fat mass, and patients with COPD present high %FM, despite not having high body weight. In addition, our data showed that more pronounced levels of obesity (i.e., morbid obesity) denote lower levels of PADL and lower exercise capacity in comparison to moderate obesity. Corroborating these results, a study by Park et al. in older women showed that those with more pronounced obesity (according to %FM) have less number of steps per day and less moderate and vigorous physical activity[33]. We believe that this is not necessarily caused only by changes due to excessive weight, but also by a reduction in the percentage of muscle mass.

It was interesting to observe that, similarly to BMI, the correlation between %FM and PADL in our sample of patients with COPD was weak. Studies concerning other populations already reported that PADL appears to be negatively associated with FM[39;40]. On the other hand, classification of obesity according to %FM yielded that all subjects with COPD in our sample were obese, which might be misinterpreted as an indication that patients with COPD do not present cachexia. It is likely that the cutoff points for %FM, which are widely used in many populations, are inappropriate for patients with COPD. In fact, FFMI value was indicative of depletion in 23 patients (32%) from the present sample ($FFMI < 16 \text{Kg/m}^2$ for men and $FFMI < 15 \text{Kg/m}^2$ for women) [41]. Therefore, both options (BMI and %FM) present clear limitations when used to reflect body composition in this population, as well as to reflect their level of physical activity. On the other hand, the

level of PADL in COPD is known to have multidimensional characteristics, being influenced by several determining factors (e.g., physical, psychosocial and environmental, and not only nutritional) [42;43]. This is likely the reason why body composition variables are not strongly related to PADL in this population, despite the fact that patients with higher degree of obesity had more marked reduction in the level of PADL.

One may argue that assessment of body composition should ideally be performed with Dual-energy X-ray absorptiometry (DEXA) and not with bioimpedance, and this can be considered as a limitation of the present study. However, electrical bioimpedance has been widely and successfully used in this population, and has provided useful and reliable information concerning body composition of patients with COPD, especially when a specific formula for these patients is used [20;44;45]. Furthermore, although the study had power enough to yield significant differences between the obese and underweight groups concerning the main outcomes (TW:87% and TPA>moderate:78%), a larger sample could have provided higher power to detect other potentially significant differences in PADL, such as between the obese and normal weight groups (see table 2). Moreover, the relatively small sample size of the present study for an investigation of such nature could lead to lack of power in certain secondary variables, such as the 6MWT (see table 2). In this sense, a larger sample could have yielded more solid results, although this would not interfere directly in the main conclusions of the study.

In conclusion, despite the high prevalence of physical inactivity in patients with COPD, those who are active in daily life have proportionally more fat free mass and less fat mass than inactive patients. More pronounced physical inactivity occurs in obese patients, although in general the correlations between body composition and PADL are weak. This suggests that body composition does not qualify as a determinant factor of the objectively assessed level of PADL in patients with COPD.

Acknowledgements

This study was supported by individual grants: FP is a researcher supported by CNPq (National Council for Scientific and Technological Development), Brazil. FM and TS were supported by MSc grants from CAPES (Brazil). CAC was supported by an MSc grant from FAPESP (Brazil). NAH was supported by a PhD grant from CAPES (Brazil).

Conflict of Interest: None

References

1. Rabe KF, Hurd S, Anzueto A, Barnes PJ, Buist SA, Calverley P, et al. (2007) Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *Am J Respir Crit Care Med* 176:532-55.
2. Landbo C, Prescott E, Lange P, Vestbo J, Almdal TP (1999) Prognostic value of nutritional status in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 160:1856-61.
3. Prescott E, Almdal T, Mikkelsen KL, Tofteng CL, Vestbo J, Lange P (2002) Prognostic value of weight change in chronic obstructive pulmonary disease: results from the Copenhagen City Heart Study. *Eur Respir J* 20:539-44.

4. Schols AM, Slangen J, Volovics L, Wouters EF (1998) Weight loss is a reversible factor in the prognosis of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 157:1791-7.
5. Slinde F, Gronberg A, Engstrom CP, Rossander-Hulthen L, Larsson S (2005) Body composition by bioelectrical impedance predicts mortality in chronic obstructive pulmonary disease patients. *Respir Med* 99:1004-9.
6. Ramachandran K, McCusker C, Connors M, Zuwallack R, Lahiri B (2008) The influence of obesity on pulmonary rehabilitation outcomes in patients with COPD. *Chron Respir Dis* 5:205-9.
7. Poulain M, Doucet M, Major GC, Drapeau V, Series F, Boulet LP, et al. (2006) The effect of obesity on chronic respiratory diseases: pathophysiology and therapeutic strategies. *CMAJ* 174:1293-9.
8. Sava F, Laviolette L, Bernard S, Breton MJ, Bourbeau J, Maltais F (2010) The impact of obesity on walking and cycling performance and response to pulmonary rehabilitation in COPD. *BMC Pulm Med* 10:55.
9. Parameswaran K, Todd DC, Soth M (2006) Altered respiratory physiology in obesity. *Can Respir J* 13:203-10.
10. Pitta F, Troosters T, Spruit MA, Probst VS, Decramer M, Gosselink R (2005) Characteristics of physical activities in daily life in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 171:972-7.
11. Hernandez NA, Teixeira DC, Probst VS, Brunetto AF, Ramos EM, Pitta F (2009) Profile of the level of physical activity in the daily lives of patients with COPD in Brazil. *J Bras Pneumol* 35:949-56.
12. Garcia-Aymerich J, Lange P, Benet M, Schnohr P, Anto JM (2006) Regular physical activity reduces hospital admission and mortality in chronic obstructive pulmonary disease: a population based cohort study. *Thorax* 61:772-8.
13. Yohannes AM, Baldwin RC, Connolly M (2002) Mortality predictors in disabling chronic obstructive pulmonary disease in old age. *Age Ageing* 31:137-40.
14. American College of Sports Medicine Position Stand. The recommended quantity and quality of exercise for developing and maintaining cardiorespiratory and muscular fitness, and flexibility in healthy adults (1998) *Med Sci Sports Exerc* 30:975-91.
15. Miller MR, Hankinson J, Brusasco V, Burgos F, Casaburi R, Coates A, et al. (2005) Standardisation of spirometry. *Eur Respir J* 26:319-38.
16. Pereira CA, Sato T, Rodrigues SC (2007) New reference values for forced spirometry in white adults in Brazil. *J Bras Pneumol* 33:397-406.
17. ATS statement: guidelines for the six-minute walk test (2002) *Am J Respir Crit Care Med* 166:1111-7.
18. Troosters T, Gosselink R, Decramer M (1999) Six minute walking distance in healthy elderly subjects. *Eur Respir J* 14:270-4.
19. Lukaski HC, Bolonchuk WW, Hall CB, Siders WA (1986) Validation of tetrapolar bioelectrical impedance method to assess human body composition. *J Appl Physiol* 60:1327-32.
20. Kyle UG, Pichard C, Rochat T, Slosman DO, Fitting JW, Thiebaud D (1998) New bioelectrical impedance formula for patients with respiratory insufficiency: comparison to dual-energy X-ray absorptiometry. *Eur Respir J* 12:960-6.
21. Pitta F, Troosters T, Spruit MA, Decramer M, Gosselink R (2005) Activity monitoring for assessment of physical activities in daily life in patients with chronic obstructive pulmonary disease. *Arch Phys Med Rehabil* 86:1979-85.
22. Langer D, Pitta F, Troosters T, Burtin C, Decramer M, Gosselink R (2009) Quantifying physical activity in COPD: different measures for different purposes. *Thorax* 64:458-9.

23. Furlanetto KC, Bisca GW, Oldemberg N, Sant'Anna TJ, Morakami FK, Camillo CA, et al. (2010) Step counting and energy expenditure estimation in patients with chronic obstructive pulmonary disease and healthy elderly: accuracy of 2 motion sensors. *Arch Phys Med Rehabil* 91:261-7.
24. Cavalheri V, Donaria L, Ferreira T, Finatti M, Camillo CA, Cipulo Ramos EM, et al. (2011) Energy expenditure during daily activities as measured by two motion sensors in patients with COPD. *Respir Med* 105:922-9.
25. Patel SA, Benzo RP, Slivka WA, Scirba FC (2007) Activity monitoring and energy expenditure in COPD patients: a validation study. *COPD* 4:107-12.
26. Nici L, Donner C, Wouters E, Zuwallack R, Ambrosino N, Bourbeau J, et al. (2006) American Thoracic Society/European Respiratory Society statement on pulmonary rehabilitation. *Am J Respir Crit Care Med* 173:1390-413.
27. Harrison B, Hubbard VS (1994) NIDDK role in obesity research. *Obes Res* 2:585-6.
28. Watz H, Waschki B, Meyer T, Magnussen H (2009) Physical activity in patients with COPD. *Eur Respir J* 33:262-72.
29. Mantoani LC, Hernandez NA, Guimaraes MM, Vitorasso RL, Probst VS, Pitta F (2011) Does the BODE index reflect the level of physical activity in daily life in patients with COPD? *Rev Bras Fisioter* 15:131-7.
30. Garcia-Aymerich J, Felez MA, Escarrabill J, Marrades RM, Morera J, Elosua R, et al. (2004) Physical activity and its determinants in severe chronic obstructive pulmonary disease. *Med Sci Sports Exerc* 36:1667-73.
31. Steuten LM, Creutzberg EC, Vrijhoef HJ, Wouters EF (2006) COPD as a multicomponent disease: inventory of dyspnoea, underweight, obesity and fat free mass depletion in primary care. *Prim Care Respir J* 15:84-91.
32. Eisner MD, Blanc PD, Sidney S, Yelin EH, Lathon PV, Katz PP, et al. (2007) Body composition and functional limitation in COPD. *Respir Res* 8:7.
33. Park J, Ishikawa-Takata K, Tanaka S, Hikiyama Y, Ohkawara K, Watanabe S, et al. (2011) Relation of body composition to daily physical activity in free-living Japanese adult women. *Br J Nutr* :1-11.
34. Puhan MA, Mador MJ, Held U, Goldstein R, Guyatt GH, Schunemann HJ (2008) Interpretation of treatment changes in 6-minute walk distance in patients with COPD. *Eur Respir J* 32:637-43.
35. Holland AE, Hill CJ, Rasekaba T, Lee A, Naughton MT, McDonald CF (2010) Updating the minimal important difference for six-minute walk distance in patients with chronic obstructive pulmonary disease. *Arch Phys Med Rehabil* 91:221-5.
36. Swinburn CR, Cooper BG, Mould H, Corris PA, Gibson GJ (1989) Adverse effect of additional weight on exercise against gravity in patients with chronic obstructive airways disease. *Thorax* 44:716-20.
37. Ofir D, Laveneziana P, Webb KA, O'Donnell DE (2007) Ventilatory and perceptual responses to cycle exercise in obese women. *J Appl Physiol* 102:2217-26.
38. Wagner PD (2008) Possible mechanisms underlying the development of cachexia in COPD. *Eur Respir J* 31:492-501.
39. Schulz LO, Schoeller DA (1994) A compilation of total daily energy expenditures and body weights in healthy adults. *Am J Clin Nutr* 60:676-81.
40. Yao M, McCrory MA, Ma G, Tucker KL, Gao S, Fuss P, et al. (2003) Relative influence of diet and physical activity on body composition in urban Chinese adults. *Am J Clin Nutr* 77:1409-16.

41. Schols AMWJ, Broekhuizen R, Weling-Scheepers CA, Wouters EF (2005) Body composition and mortality in chronic obstructive pulmonary disease. *Am J Clin Nutr* 82:53-59.
42. Caspersen CJ, Powell KE, Christenson GM (1985) Physical activity, exercise and physical fitness: definitions and distinctions for health-related research. *Public Health Rep* 100:126-131.
43. van Sluijs EM, Griffin SJ, van Poppel MN (2007) A cross-sectional study of awareness of physical activity: associations with personal, behavioral and psychosocial factors. *Int J Behav Nutr Phys Act* 4:53.
44. Rutten EP, Spruit MA, Wouters EF (2010) Critical view on diagnosing muscle wasting by single-frequency bio-electrical impedance in COPD. *Respir Med* 104:91-8.
45. Schols AM, Wouters EF, Soeters PB, Westerterp KR (1991) Body composition by bioelectrical-impedance analysis compared with deuterium dilution and skinfold anthropometry in patients with chronic obstructive pulmonary disease. *Am J Clin Nutr* 53:421-4.

TABELAS

Table 1 – Results from the general group of patients with Chronic Obstructive Pulmonary Disease (COPD) and the physically active and inactive subgroups according to American College of Sports Medicine (ACSM) minimum recommendations of daily physical activity.

Characteristics	Total n=74	Active n=22	Inactive n=52	p (act vs inact)
Age (years)	65±9	65±12	65±7	0.90
Gender (M/F)	45/29	15/7	30/22	0.86
Weight (Kg)	69±16	61±12	73±16	0.002
Height (cm)	160±9	161±10	160±7	0.82
FEV ₁ (%pred)	40±15	44±15	39±14	0.18
BMI (Kg/m ²)	27±6	23±4	28±6	0.001
FFMI (Kg/m ²)	18±3	16±2	18±4	0.009
FFM (Kg)	46±11	42±11	47±10	0.05
FM (Kg)	23±8	18±6	25±8	0.0005
FFM (% of BW)	67±8	70±8	65±7	0.03
FM (% of BW)	33±8	30±8	35±7	0.03
Metabolic rate (Kcal)	1396±322	1285±308	1442±319	0.05
6MWT (m)	428±94	454±113	416±84	0.04
6MWT (%pred)	72±16	74±18	71±15	0.21

Subgroup Active: achieves at least 30 minutes per day of physical activity of at least moderate intensity, in accordance with the ACSM recommendations; Subgroup Inactive: does not achieve these minimum recommendations. Data are expressed as mean (standard deviation). FEV₁ = forced expiratory volume in the first second; BMI = body mass index; FFMI = fat free mass index; FFM = fat free mass; FM = fat mass; BW = body weight; 6MWT = 6-minute walking test.

Table 2 – Comparison of patients with Chronic Obstructive Pulmonary Disease (COPD) classified according to the body composition based on the body mass index (BMI)

	Underweight n=14	Normal weight n=17	Overweight n=22	Obese n=21
Age (years)	65±11	66±11	66±8	65±6
Gender (M/F)	10/4	11/7	14/8	10/11
BMI (Kg.m ⁻²)	19±1	23±1*	28±1 ^o	35±4 [#]
FEV ₁ (%pred)	36±14	35±15	40±14	47±13 *
6MWT (m)	460±61	447±95	410±115	406±83
TPA>mod (min/day)	42±43	31±33	23±36	11±12 *
TW (min/day)	73±31	69±34	53±30	46±26 * +

Data are expressed as mean ± standard deviation. FEV₁ = forced expiratory volume in the first second; 6MWT = 6-minute walking test; TPA>mod = time spent per day in physical activities of at least moderate intensity; TW = time spent walking per day. Underweight: BMI<21 Kg.m⁻²; Normal weight: 21≤BMI<25Kg.m⁻²; Overweight: 25≤BMI<30Kg.m⁻² and Obese: BMI>30Kg.m⁻².

* p <0.05 versus underweight

^o p <0.05 versus underweight and normal weight

[#] p <0.05 versus underweight, normal weight and overweight

+ p <0.05 versus normal weight

Table 3 – Comparison of patients with Chronic Obstructive Pulmonary Disease (COPD) classified according to the body composition based on the percentage of fat mass (FM%).

	Moderate Obesity n=11	High Obesity n=30	Morbid Obesity n=30
Age (years)	65±10	62±7	69±8 ⁺
Gender (M/F)	7/4	15/15	20/10
FEV ₁ (%pred)	35±14	44±14	38±14
6MWT(m)	475±75	445±88	389±98*
TPA>mod (min/day)	47±45	19±20 *	20±26 *
TW (min/day)	72±36	62±29	43±27 * +

Data are expressed as mean ± standard deviation. FEV₁ = forced expiratory volume in the first second; 6MWT = 6-minute walking test; TPA>mod = time spent per day in physical activities of at least moderate intensity; TW = time spent walking per day. Moderate obesity: Women: 30≤%FM<35 - Men: 20≤%FM<25; high obesity: W: 35≤%FM<40 - M: 25≤%FM<30 and morbid obesity: W: %FM≥40 - M: %FM≥30.

* p<0.05 versus Moderate Obesity

+ p <0.05 versus High Obesity

4 CONCLUSÕES GERAIS

Pacientes com DPOC, que são fisicamente ativos na vida diária têm proporcionalmente mais massa magra e menos massa de gordura do que os pacientes fisicamente inativos. Maiores níveis de inatividade física ocorrem em pacientes obesos, embora em geral as correlações entre a composição corporal e AFVD sejam fracas. Isto sugere que a apesar da composição corporal de pacientes ativos ser diferente da dos inativos e de pacientes obesos possuírem pior AFVD, a composição corporal de forma geral não se qualifica como fator determinante do nível de AFVD avaliado objetivamente em pacientes com DPOC. Estes resultados nos levam a crer que, apesar da literatura sugerir que a obesidade é protetora em pacientes com DPOC, atenção especial deve ser dada ao nível de atividade de física de pacientes obesos, já que estes são os principais acometidos pela inatividade física.

REFERÊNCIAS

- (1) Rabe KF, Hurd S, Anzueto A, Barnes PJ, Buist SA, Calverley P, et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *Am J Respir Crit Care Med* 2007 Sep 15;176(6):532-55.
- (2) Landbo C, Prescott E, Lange P, Vestbo J, Almdal TP. Prognostic value of nutritional status in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1999 Dec;160(6):1856-61.
- (3) Slinde F, Gronberg A, Engstrom CP, Rossander-Hulthen L, Larsson S. Body composition by bioelectrical impedance predicts mortality in chronic obstructive pulmonary disease patients. *Respir Med* 2005 Aug;99(8):1004-9.
- (4) Prescott E, Almdal T, Mikkelsen KL, Tofteng CL, Vestbo J, Lange P. Prognostic value of weight change in chronic obstructive pulmonary disease: results from the Copenhagen City Heart Study. *Eur Respir J* 2002 Sep;20(3):539-44.
- (5) Ramachandran K, McCusker C, Connors M, Zuwallack R, Lahiri B. The influence of obesity on pulmonary rehabilitation outcomes in patients with COPD. *Chron Respir Dis* 2008;5(4):205-9.
- (6) Poulain M, Doucet M, Major GC, Drapeau V, Series F, Boulet LP, et al. The effect of obesity on chronic respiratory diseases: pathophysiology and therapeutic strategies. *CMAJ* 2006 Apr 25;174(9):1293-9.
- (7) Parameswaran K, Todd DC, Soth M. Altered respiratory physiology in obesity. *Can Respir J* 2006 May;13(4):203-10.
- (8) Sava F, Laviolette L, Bernard S, Breton MJ, Bourbeau J, Maltais F. The impact of obesity on walking and cycling performance and response to pulmonary rehabilitation in COPD. *BMC Pulm Med* 2010;10:55.
- (9) American College of Sports Medicine Position Stand. The recommended quantity and quality of exercise for developing and maintaining cardiorespiratory and muscular fitness, and flexibility in healthy adults. *Med Sci Sports Exerc* 1998 Jun;30(6):975-91.
- (10) Pitta F, Troosters T, Probst VS, Langer D, Decramer M, Gosselink R. Are patients with COPD more active after pulmonary rehabilitation? *Chest* 2008 Aug;134(2):273-80.
- (11) Park J, Ishikawa-Takata K, Tanaka S, Hikiyama Y, Ohkawara K, Watanabe S, et al. Relation of body composition to daily physical activity in free-living Japanese adult women. *Br J Nutr* 2011 May 17;1-11.
- (12) World Health Report. Primary Health Care, WHO Library Cataloguing . 2008. www.who.int/en. Último acesso em setembro de 2011.

- (13) Qaseem A, Wilt TJ, Weinberger SE, Hanania NA, Criner G, van der Molen T, et al. Diagnosis and Management of Stable Chronic Obstructive Pulmonary Disease: A Clinical Practice Guideline Update from the American College of Physicians, American College of Chest Physicians, American Thoracic Society, and European Respiratory Society. *Ann Intern Med* 2011 Aug 2;155(3):179-91.
- (14) Murray CJ, Lopez AD. Alternative projections of mortality and disability by cause 1990-2020: Global Burden of Disease Study. *Lancet* 1997 May 24;349(9064):1498-504.
- (15) Sullivan SD, Ramsey SD, Lee TA. The economic burden of COPD. *Chest* 2000 Feb;117(2 Suppl):5S-9S.
- (16) Soriano JB, Maier WC, Egger P, Visick G, Thakrar B, Sykes J, et al. Recent trends in physician diagnosed COPD in women and men in the UK. *Thorax* 2000 Sep;55(9):789-94.
- (17) Campos H. Doença pulmonar obstrutiva crônica. *J Bras Med* 1993;64(6):98-114.
- (18) Menezes AM. Epidemiologia da bronquite crônica e do enfisema (DPOC): até onde sabemos? *J Pneumol* 1997;23(3):153-7.
- (19) Mantoani LC, Hernandez NA, Guimaraes MM, Vitorasso RL, Probst VS, Pitta F. Does the BODE index reflect the level of physical activity in daily life in patients with COPD? *Rev Bras Fisioter* 2011 Apr;15(2):131-7.
- (20) Troosters T, Gosselink R, Decramer M. Pulmonary rehabilitation in patients with severe chronic obstructive pulmonary disease. *Monaldi Arch Chest Dis* 1999 Dec;54(6):510-3.
- (21) Hernandez NA, Teixeira DC, Probst VS, Brunetto AF, Ramos EM, Pitta F. Profile of the level of physical activity in the daily lives of patients with COPD in Brazil. *J Bras Pneumol* 2009 Oct;35(10):949-56.
- (22) Paz-Diaz H, Montes de OM, Lopez JM, Celli BR. Pulmonary rehabilitation improves depression, anxiety, dyspnea and health status in patients with COPD. *Am J Phys Med Rehabil* 2007 Jan;86(1):30-6.
- (23) Langer D, Hendriks E, Burtin C, Probst V, van der Schans C, Paterson W, et al. A clinical practice guideline for physiotherapists treating patients with chronic obstructive pulmonary disease based on a systematic review of available evidence. *Clin Rehabil* 2009 May;23(5):445-62.
- (24) World Health Organization. WHO Global Database on Body Mass Index (BMI) Obesity and overweight . 2011. www.who.int/mediacentre/factsheets/fs311. Último acesso em setembro de 2011.
- (25) Eckersley RM. Losing the battle of the bulge: causes and consequences of increasing obesity. *Med J Aust* 2001 Jun 4;174(11):590-2.

- (26) World Health Organization. Global database on body mass index. 2011. <<http://apps.who.int/bmi/index.jsp>>. Último acesso em setembro de 2011.
- (27) Visscher TL, Seidell JC, Menotti A, Blackburn H, Nissinen A, Feskens EJ, et al. Underweight and overweight in relation to mortality among men aged 40-59 and 50-69 years: the Seven Countries Study. *Am J Epidemiol* 2000 Apr 1;151(7):660-6.
- (28) Obesity: preventing and managing the global epidemic. Report of a WHO consultation. *World Health Organ Tech Rep Ser* 2000;894:i-253.
- (29) Geloneze B, Mancini MC, Coutinho W. Obesity: knowledge, care, and commitment, but not yet cure. *Arq Bras Endocrinol Metabol* 2009 Mar;53(2):117-9.
- (30) Overweight, obesity, and health risk. National Task Force on the Prevention and Treatment of Obesity. *Arch Intern Med* 2000 Apr 10;160(7):898-904.
- (31) Naimark A, Cherniack RM. Compliance of the respiratory system and its components in health and obesity. *J Appl Physiol* 1960 May;15:377-82.
- (32) Sin DD, Jones RL, Man SF. Obesity is a risk factor for dyspnea but not for airflow obstruction. *Arch Intern Med* 2002 Jul 8;162(13):1477-81.
- (33) Calle EE, Thun MJ, Petrelli JM, Rodriguez C, Heath CW, Jr. Body-mass index and mortality in a prospective cohort of U.S. adults. *N Engl J Med* 1999 Oct 7;341(15):1097-105.
- (34) Okorodudu DO, Jumean MF, Montori VM, Romero-Corral A, Somers VK, Erwin PJ, et al. Diagnostic performance of body mass index to identify obesity as defined by body adiposity: a systematic review and meta-analysis. *Int J Obes (Lond)* 2010 May;34(5):791-9.
- (35) Adams TD, Heath EM, LaMonte MJ, Gress RE, Pendleton R, Strong M, et al. The relationship between body mass index and per cent body fat in the severely obese. *Diabetes Obes Metab* 2007 Jul;9(4):498-505.
- (36) Jackson AS, Ellis KJ, McFarlin BK, Sailors MH, Bray MS. Body mass index bias in defining obesity of diverse young adults: the Training Intervention and Genetics of Exercise Response (TIGER) study. *Br J Nutr* 2009 Oct;102(7):1084-90.
- (37) Caspersen CJ, Powell KE, Christenson GM. Physical activity, exercise, and physical fitness: definitions and distinctions for health-related research. *Public Health Rep* 1985 Mar;100(2):126-31.
- (38) World Health Organization. Physical activity: direct and indirect health benefits. 2002. www.who.int/topics/physical_activity. Último acesso em setembro de 2011.
- (39) Physical activity programs and behavior counseling in older adult populations. *Med Sci Sports Exerc* 2004 Nov;36(11):1997-2003.

- (40) Shaaban R, Leynaert B, Soussan D, Anto JM, Chinn S, de MR, et al. Physical activity and bronchial hyperresponsiveness: European Community Respiratory Health Survey II. *Thorax* 2007 May;62(5):403-10.
- (41) CENTERS FOR DISEASE CONTROL AND PREVENTION. Improving nutrition and increasing physical activity. 2009. www.cdc.gov/physicalactivity. Último acesso em setembro de 2011.
- (42) World Health Organization. Sedentary lifestyle: a global public health problem. 2002. www.who.int/dietphysicalactivity. Último acesso em setembro de 2011.
- (43) World Health Organization. Global Health Observatory (GHO): Prevalence of insufficient physical activity. 2011. www.who.int/dietphysicalactivity. Último acesso em setembro de 2011.
- (44) Pitta F, Troosters T, Probst VS, Spruit MA, Decramer M, Gosselink R. Quantifying physical activity in daily life with questionnaires and motion sensors in COPD. *Eur Respir J* 2006 May;27(5):1040-55.
- (45) Manini TM, Everhart JE, Patel KV, Schoeller DA, Colbert LH, Visser M, et al. Daily activity energy expenditure and mortality among older adults. *JAMA* 2006 Jul 12;296(2):171-9.
- (46) Steele BG, Holt L, Belza B, Ferris S, Lakshminaryan S, Buchner DM. Quantitating physical activity in COPD using a triaxial accelerometer. *Chest* 2000 May;117(5):1359-67.
- (47) Pitta F, Troosters T, Spruit MA, Decramer M, Gosselink R. Activity monitoring for assessment of physical activities in daily life in patients with chronic obstructive pulmonary disease. *Arch Phys Med Rehabil* 2005 Oct;86(10):1979-85.
- (48) Furlanetto KC, Bisca GW, Oldenberg N, Sant'Anna TJ, Morakami FK, Camillo CA, et al. Step counting and energy expenditure estimation in patients with chronic obstructive pulmonary disease and healthy elderly: accuracy of 2 motion sensors. *Arch Phys Med Rehabil* 2010 Feb;91(2):261-7.
- (49) Patel SA, Benzo RP, Slivka WA, Sciurba FC. Activity monitoring and energy expenditure in COPD patients: a validation study. *COPD* 2007 Jun;4(2):107-12.
- (50) ten Hacken NH. Physical inactivity and obesity: relation to asthma and chronic obstructive pulmonary disease? *Proc Am Thorac Soc* 2009 Dec;6(8):663-7.
- (51) Schols AM, Slangen J, Volovics L, Wouters EF. Weight loss is a reversible factor in the prognosis of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1998 Jun;157(6 Pt 1):1791-7.
- (52) Garcia-Aymerich J, Lange P, Benet M, Schnohr P, Anto JM. Regular physical activity reduces hospital admission and mortality in chronic obstructive pulmonary disease: a population based cohort study. *Thorax* 2006 Sep;61(9):772-8.

- (53) Yohannes AM, Baldwin RC, Connolly M. Mortality predictors in disabling chronic obstructive pulmonary disease in old age. *Age Ageing* 2002 Mar;31(2):137-40.
- (54) Weiner P, Waizman J, Weiner M, Rabner M, Magadle R, Zamir D. Influence of excessive weight loss after gastroplasty for morbid obesity on respiratory muscle performance. *Thorax* 1998 Jan;53(1):39-42.
- (55) Chlif M, Keochkerian D, Mourlhon C, Choquet D, Ahmaidi S. Noninvasive assessment of the tension-time index of inspiratory muscles at rest in obese male subjects. *Int J Obes (Lond)* 2005 Dec;29(12):1478-83.
- (56) Seres L, Lopez-Ayerbe J, Coll R, Rodriguez O, Manresa JM, Marrugat J, et al. [Cardiopulmonary function and exercise capacity in patients with morbid obesity]. *Rev Esp Cardiol* 2003 Jun;56(6):594-600.

ANEXOS

ANEXO A

Normas de formatação do periódico LUNG

<http://www.springer.com/medicine/internal/journal/408>

Instructions for Authors

1. Types of papers

Original Article, Original Research, State of the Art Review, Brief Report, Editorial, Book Review, and Lung Images

2. Editorial procedure

Additional Info: It would be useful if the authors would submit the names, telephone and fax numbers, and addresses of three to five people who might review the manuscript.

3. Manuscript submission

Manuscript Submission

Submission of a manuscript implies: that the work described has not been published before; that it is not under consideration for publication anywhere else; that its publication has been approved by all co-authors, if any, as well as by the responsible authorities – tacitly or explicitly – at the institute where the work has been carried out. The publisher will not be held legally responsible should there be any claims for compensation.

Permissions

Authors wishing to include figures, tables, or text passages that have already been published elsewhere are required to obtain permission from the copyright owner(s) for both the print and online format and to include evidence that such permission has been granted when submitting their papers. Any material received without such evidence will be assumed to originate from the authors.

Online Submission

Authors should submit their manuscripts online. Electronic submission substantially reduces the editorial processing and reviewing times and shortens overall publication times. Please follow the hyperlink “Submit online” on the

right and upload all of your manuscript files following the instructions given on the screen.

4. Title page

The title page should include:

- The name(s) of the author(s)
- A concise and informative title
- The affiliation(s) and address(es) of the author(s)
- The e-mail address, telephone and fax numbers of the

corresponding author

Abstract

Please provide a structured abstract of 150 to 250 words which should be divided into the following sections:

- Purpose (stating the main purposes and research question)
- Methods
- Results
- Conclusions

Keywords

Please provide 4 to 6 keywords which can be used for indexing purposes.

5. Text

Text Formatting

Manuscripts should be submitted in Word.

- Use a normal, plain font (e.g., 10-point Times Roman) for text.
- Use italics for emphasis.
- Use the automatic page numbering function to number the pages.
- Do not use field functions.
- Use tab stops or other commands for indents, not the space bar.
- Use the table function, not spreadsheets, to make tables.

- Use the equation editor or MathType for equations.

Note: If you use Word 2007, do not create the equations with the default equation editor but use the Microsoft equation editor or MathType instead.

- Save your file in doc format. Do not submit docx files.
- Word template (zip, 154 kB)

Manuscripts with mathematical content can also be submitted in LaTeX.

- LaTeX macro package (zip, 182 kB)

Headings

Please use no more than three levels of displayed headings.

Abbreviations

Abbreviations should be defined at first mention and used consistently thereafter.

Footnotes

Footnotes can be used to give additional information, which may include the citation of a reference included in the reference list. They should not consist solely of a reference citation, and they should never include the bibliographic details of a reference. They should also not contain any figures or tables.

Footnotes to the text are numbered consecutively; those to tables should be indicated by superscript lower-case letters (or asterisks for significance values and other statistical data). Footnotes to the title or the authors of the article are not given reference symbols.

Always use footnotes instead of endnotes.

Acknowledgments

Acknowledgments of people, grants, funds, etc. should be placed in a separate section before the reference list. The names of funding organizations should be written in full.

6. Letters to the Editor

Text is limited to 750 words, with no abstract. There may be 1 figure, up to 3 references, and no more than 3 authors, with author affiliations only including main institution, place name and (state plus) country (i.e. no departments, etc.).

7. Scientific style

- Please always use internationally accepted signs and symbols for units (SI units).

- Nomenclature: Insofar as possible, authors should use systematic names similar to those used by Chemical Abstract Service or IUPAC.

- Please use the standard mathematical notation for formulae, symbols etc.:

Italic for single letters that denote mathematical constants, variables, and unknown quantities

Roman/upright for numerals, operators, and punctuation, and commonly defined functions or abbreviations, e.g., cos, det, e or exp, lim, log, max, min, sin, tan, d (for derivative)

Bold for vectors, tensors, and matrices.

8. Terminology

- Generic names of drugs and pesticides are preferred; if trade names are used, the generic name should be given at first mention.

9. References

- Citation

Reference citations in the text should be identified by numbers in square brackets. Some examples:

1. Negotiation research spans many disciplines [3].
2. This result was later contradicted by Becker and Seligman [5].
3. This effect has been widely studied [1-3, 7].

- Reference list

The list of references should only include works that are cited in the text and that have been published or accepted for publication. Personal

communications and unpublished works should only be mentioned in the text. Do not use footnotes or endnotes as a substitute for a reference list.

The entries in the list should be numbered consecutively.

- Journal article

Gamelin FX, Baquet G, Berthoin S, Thevenet D, Nourry C, Nottin S, Bosquet L (2009) Effect of high intensity intermittent training on heart rate variability in prepubescent children. *Eur J Appl Physiol* 105:731-738. doi: 10.1007/s00421-008-0955-8

Ideally, the names of all authors should be provided, but the usage of “et al” in long author lists will also be accepted:

Smith J, Jones M Jr, Houghton L et al (1999) Future of health insurance. *N Engl J Med* 965:325–329

- Article by DOI

Slifka MK, Whitton JL (2000) Clinical implications of dysregulated cytokine production. *J Mol Med*. doi:10.1007/s001090000086

- Book

South J, Blass B (2001) *The future of modern genomics*. Blackwell, London

- Book chapter

Brown B, Aaron M (2001) The politics of nature. In: Smith J (ed) *The rise of modern genomics*, 3rd edn. Wiley, New York, pp 230-257

- Online document

Cartwright J (2007) Big stars have weather too. IOP Publishing PhysicsWeb. <http://physicsweb.org/articles/news/11/6/16/1>. Accessed 26 June 2007

- Dissertation

Trent JW (1975) *Experimental acute renal failure*. Dissertation, University of California

Always use the standard abbreviation of a journal’s name according to the ISSN List of Title Word Abbreviations, see

- www.issn.org/2-22661-LTWA-online.php

For authors using EndNote, Springer provides an output style that supports the formatting of in-text citations and reference list.

- EndNote style (zip, 2 kB)

Authors preparing their manuscript in LaTeX can use the bibtex file `spbasic.bst` which is included in Springer's LaTeX macro package.

10. Tables

- All tables are to be numbered using Arabic numerals.
- Tables should always be cited in text in consecutive numerical order.
 - For each table, please supply a table caption (title) explaining the components of the table.
 - Identify any previously published material by giving the original source in the form of a reference at the end of the table caption.
 - Footnotes to tables should be indicated by superscript lower-case letters (or asterisks for significance values and other statistical data) and included beneath the table body.

11. Artwork

For the best quality final product, it is highly recommended that you submit all of your artwork – photographs, line drawings, etc. – in an electronic format. Your art will then be produced to the highest standards with the greatest accuracy to detail. The published work will directly reflect the quality of the artwork provided.

12. Integrity of research and reporting

Ethical standards

Manuscripts submitted for publication must contain a declaration that the experiments comply with the current laws of the country in which they were performed. Please include this note in a separate section before the reference list.

Conflict of interest

All benefits in any form from a commercial party related directly or indirectly to the subject of this manuscript or any of the authors must be acknowledged. For each source of funds, both the research funder and the grant number should be given. This note should be added in a separate section before the reference list.

If no conflict exists, authors should state: The authors declare that they have no conflict of interest.